



AD 711804


THEMIS SIGNAL ANALYSIS STATISTICS RESEARCH PROGRAM

CORRELATION BETWEEN TWO HOTELLING'S  $T^2$

by

A. M. Kshirsagar\* and John C. Young

Technical Report No. 79  
Department of Statistics THEMIS Contract



Reproduction in whole or in part is permitted  
for any purpose of the United States Government.

This document has been approved for public release  
and sale; its distribution is unlimited.

**Department of Statistics**  
**Southern Methodist University**  
Dallas, Texas 75222

THEMIS SIGNAL ANALYSIS STATISTICS RESEARCH PROGRAM

CORRELATION BETWEEN TWO HOTELLING'S  $T^2$

by

A. M. Kshirsagar and John C. Young

Technical Report No. 79  
Department of Statistics THEMIS Contract

August 1, 1970

Research sponsored by the Office of Naval Research  
Contract N00014-68-A-0515  
Project NR 042-260

Reproduction in whole or in part is permitted  
for any purpose of the United States Government.

This document has been approved for public release  
and sale; its distribution is unlimited.

DEPARTMENT OF STATISTICS  
Southern Methodist University

# CORRELATION BETWEEN TWO HOTELLING'S $T^2$

by

A. M. Kshirsagar\* and John C. Young

Southern Methodist University  
Dallas, Texas 75222

## 1. Introduction:

Let  $B$  be a  $p \times p$  symmetric matrix having the Wishart distribution

$$(1.1) \quad W_p(B|I|f)dB = C_{pf}|B|^{(f-p-1)/2} e^{-1/2 \text{tr} B} dB,$$

where

$$(1.2) \quad C_{pf}^{-1} = 2^{fp/2} \pi^{p(p-1)/4} \prod_{i=1}^p \Gamma\left(\frac{f+1-i}{2}\right),$$

and  $dB$  stands for the product of the differentials of the  $p(p+1)/2$  distinct elements of  $B$ . Let  $\underline{x}$  and  $\underline{y}$  be two vector variables of  $p$  components, distributed independently of  $B$ , and also independently of each other, as

$$(1.3) \quad \frac{1}{(2\pi)^{p/2}} e^{-1/2 \underline{x}'\underline{x}} d\underline{x},$$

and

$$(1.4) \quad \frac{1}{(2\pi)^{p/2}} e^{-1/2 \underline{y}'\underline{y}} d\underline{y}$$

respectively. While considering the problem of multivariate statistical outliers, Wilks (1963) used statistics of the type,

---

\*Research supported by ONR Contract N00014-68-A-0515, Department of Statistics THEMIS Project.

$$(1.5) \quad r = |B + \underline{y}\underline{y}'| / |B + \underline{x}\underline{x}' + \underline{y}\underline{y}'|, \quad s = |B + \underline{x}\underline{x}'| / |B + \underline{x}\underline{x}' + \underline{y}\underline{y}'|.$$

He has remarked that the exact distribution (joint) of  $r$  and  $s$  is complicated and has given the expected values, variances and covariance of  $r$  and  $s$ . Unfortunately, his expressions for the variance and covariance are in error. The purpose of this note is to derive the exact joint distribution of  $r$  and  $s$  and to give correct expressions for the moments.

## 2. Joint distribution:

In the joint distribution of  $B$ ,  $\underline{x}$  and  $\underline{y}$ , make the following transformation

$$(2.1) \quad \begin{aligned} A &= B + \underline{x}\underline{x}' + \underline{y}\underline{y}', \\ \underline{u} &= A^{-1/2} \underline{x}, \\ \underline{v} &= A^{-1/2} \underline{y}, \end{aligned}$$

where  $A^{-1/2}$  is any matrix such that  $A^{-1/2} \cdot A^{-1/2} = A^{-1}$ . The Jacobian of transformation from  $B$  to  $A$  is 1 and that from  $\underline{x}$  to  $\underline{u}$  or  $\underline{y}$  to  $\underline{v}$  is  $|A|^{1/2}$  and hence, the joint distribution of  $A$ ,  $\underline{u}$  and  $\underline{v}$  comes out as

$$(2.2) \quad \frac{C_{pf}}{(2\pi)^p} |A|^{\frac{(f+2)-p-1}{2}} e^{-1/2 \operatorname{tr} A} \cdot |I - \underline{u}\underline{u}' - \underline{v}\underline{v}'|^{\frac{f-p-1}{2}} dA d\underline{u} d\underline{v},$$

$$\text{as } |B| = |A - A^{1/2} \underline{u}\underline{u}' A^{1/2} - A^{1/2} \underline{v}\underline{v}' A^{1/2}| = |A| |I - \underline{u}\underline{u}' - \underline{v}\underline{v}'|.$$

This shows that  $A$  has a Wishart distribution of  $f+2$  degrees of freedom and is independent of  $\underline{u}$  and  $\underline{v}$ . Splitting the constant suitably, the joint distribution of  $\underline{u}$  and  $\underline{v}$  is

$$(2.3) \quad \frac{\Gamma(f+1)}{(2\pi)^p \Gamma(f-p+1)} |I - \underline{u}\underline{u}' - \underline{v}\underline{v}'|^{\frac{f-p-1}{2}} d\underline{u} d\underline{v}.$$

Observe that the statistics  $r$ ,  $s$  of Wilks are given by

$$(2.4) \quad r = \frac{|B + \underline{y}\underline{y}'|}{|B + \underline{x}\underline{x}' + \underline{y}\underline{y}'|} = \frac{|A - \underline{x}\underline{x}'|}{|A|} = |I - \underline{u}\underline{u}'| = 1 - \underline{u}'\underline{u},$$

and

$$(2.5) \quad s = \frac{|B+xx'|}{|B+xx'+yy'|} = 1 - \underline{v}'\underline{v} \quad .$$

Also observe that in (2.3)

$$(2.6) \quad \begin{aligned} |I - \underline{u}\underline{u}' - \underline{v}\underline{v}'| &= (1 - \underline{u}'\underline{u})(1 - \underline{v}'\underline{v}) - (\underline{u}'\underline{v})^2 \\ &= rs - (\underline{u}'\underline{v})^2 \quad . \end{aligned}$$

In (2.3), transform from  $\underline{v}$  to  $\underline{w} = [w_1, w_2, \dots, w_p]'$ , by an orthogonal transformation

$$(2.7) \quad \underline{w} = L\underline{v} \quad ,$$

where

$L$  is a  $p \times p$  orthogonal matrix, whose last row is  $\underline{u}'/\sqrt{\underline{u}'\underline{u}}$ . The Jacobian of this transformation is  $|L| = 1$  and  $\underline{v}'\underline{v} = \underline{w}'\underline{w} = 1-s$ . Also

$$(2.8) \quad \underline{u}'\underline{v} = \underline{u}'L'L\underline{v} = [0 \dots 0, \sqrt{\underline{u}'\underline{u}}] \underline{w} = \sqrt{1-r} \cdot w_p \quad .$$

The joint distribution of  $\underline{u}$  and  $\underline{w}$  is, therefore,

$$(2.9) \quad \frac{\Gamma(f+1)}{(2\pi)^p \Gamma(f-p+1)} \left\{ rs - (1-r)w_p^2 \right\}^{\frac{f-p-1}{2}} d\underline{u}d\underline{w} \quad .$$

From  $\underline{u}$ , transform to  $r = 1 - \underline{u}'\underline{u}$  and  $p-1$  other variables

$\phi_1, \phi_2, \dots, \phi_{p-1}$  by

$$(2.10) \quad \begin{aligned} u_1 &= (1-r)^{1/2} \cos\phi_1 \cos\phi_2 \dots \cos\phi_{p-1} \quad , \\ u_j &= (1-r)^{1/2} \cos\phi_1 \cos\phi_2 \dots \cos\phi_{p-j} \sin\phi_{p-j+1} \\ &\quad (j=2, 3, \dots, p) \end{aligned}$$

Similarly, transform from  $\underline{w}$  to  $s = 1 - \underline{w}'\underline{w}$  and  $p-1$  other variables

$\theta_1, \theta_2, \dots, \theta_{p-1}$  by

$$(2.11) \quad \begin{aligned} v_1 &= (1-s)^{1/2} \cos\theta_1 \cos\theta_2 \dots \cos\theta_{p-1} \quad , \\ v_j &= (1-s)^{1/2} \cos\theta_1 \cos\theta_2 \dots \cos\theta_{p-j} \sin\theta_{p-j+1} \quad . \\ &\quad (j=2, 3, \dots, p) \end{aligned}$$

The Jacobian of transformation from  $\underline{u}$  to  $r, \phi_1, \dots, \phi_{p-1}$  is

$$\frac{1}{2}(1-r)^{\frac{1}{2}p-1} \prod_{i=1}^{p-2} \cos \phi_i^{p-i-1}$$

and a similar expression in  $s$  and  $\theta_1$  for the Jacobian of transformation from  $\underline{w}$  to  $s$  and the  $\theta$ 's. Now  $\theta_{p-1}$  and  $\phi_{p-1}$  vary from 0 to  $2\pi$ , the other  $\theta$ 's and  $\phi$ 's vary from  $-\pi/2$  to  $\pi/2$  while  $r$  and  $s$  vary from 0 to 1. Integrating out all the  $\phi$ 's and all  $\theta$ 's except  $\theta_1$ , we obtain the joint distribution of  $r, s$  and  $\theta_1$  as

$$(2.12) \quad \frac{\Gamma(f+1)}{4\pi\Gamma(p-1)\Gamma(f-p+1)} \{rs - (1-r)(1-s)\sin^2\theta\}^{\frac{f-p-1}{2}} \cos^{p-2}\theta \, dr ds d\theta$$

where  $\theta_1$  is replaced by  $\theta$ .

The joint distribution of  $r, s$  alone can now be obtained by integrating out  $\theta$  but this does not seem to yield a manageable expression, as the bracket in (2.12) will have to be expanded in a series.

### 3. Moments of $r, s$ .

Only the product moment of  $r$  and  $s$  is difficult to obtain. The mean and variance of  $r$  (or  $s$ ) can be very easily obtained from the marginal distribution of  $r$ , which is related to the well-known Hotelling's

$T^2$  by  $r = \frac{1}{1 + \left(\frac{T^2}{f+1}\right)}$ . In the joint distribution of  $\underline{u}$  and  $\underline{v}$ , given by

(2.3), if we transform to  $\underline{z} = [z_1, \dots, z_p]'$  from  $\underline{v}$  by

$$(3.1) \quad \underline{v} = (\mathbf{I} - \underline{u}\underline{u}')^{1/2} \underline{z},$$

we shall find that  $\underline{u}$  and  $\underline{z}$  are independently distributed as

$$(3.2) \quad K(\underline{u}|f) d\underline{u} = \frac{f}{\pi^{p/2}(f-p)} \cdot \frac{\Gamma(f/2)}{\frac{\pi}{2}(f-p)} |\mathbf{I} - \underline{u}\underline{u}'|^{\frac{f-p}{2}} d\underline{u}$$

$$(3.3) \quad \text{and } K(\underline{z}|f-1) d\underline{z}, \text{ respectively.}$$

From (3.2), one can easily show that

$$(3.4) \quad E(r^h) = E(1 - \underline{u}'\underline{u})^h = E|I - \underline{u}\underline{u}'|^h \\ = \frac{f(f+2h-p)}{(f-p)(f+2h)} \cdot \frac{\Gamma(\frac{f-p}{2} + h)\Gamma(\frac{f}{2})}{\Gamma(\frac{f}{2} + h)\Gamma(\frac{f-p}{2})}$$

This will also be the  $h^{\text{th}}$  moment of  $s$  by symmetry. This leads to

$$(3.5) \quad E(r) = \frac{f-p+2}{f+2}, \quad v(r) = \frac{2p(f-p+2)}{(f+2)^2(f+4)}.$$

$$\text{Now } \text{Cov}(r, s) = E\{(1 - \underline{u}'\underline{u})(1 - \underline{v}'\underline{v})\} - E(r)E(s)$$

$$= E\{(1 - \underline{u}'\underline{u})[1 - \underline{z}(I - \underline{u}\underline{u}')\underline{z}]\} - \{E(r)\}^2 \quad \text{by (3.1)}$$

$$= E(r) - E\{r[\underline{z}'\underline{z} - (\underline{z}'\underline{u})^2]\} - \{E(r)\}^2$$

$$(3.6) \quad = E(r) - E(r)E(\underline{z}'\underline{z}) + E\{r(\underline{z}'\underline{u})^2\} - \{E(r)\}^2,$$

as  $\underline{z}$  and  $r$  are independent. Since  $\underline{z}$  has the same distribution as  $\underline{u}$  with  $f$  changed  $f-1$ ,

$$(3.7) \quad E(\underline{z}'\underline{z}) = 1 - E(1 - \underline{u}'\underline{u}) \text{ with } f \text{ replaced by } f-1 \\ = \frac{p}{f+1}$$

Hence (3.6) reduces to

$$(3.8) \quad \text{Cov}(r, s) = \frac{-p(f-p+2)}{(f+1)(f+2)^2} + E\{r(\underline{z}'\underline{u})^2\}.$$

Now

$$(3.9) \quad E\{r(\underline{z}'\underline{u})^2\} = \int (1 - \underline{u}'\underline{u}) (\underline{z}'\underline{u})^2 K(\underline{u}|f) K(\underline{z}|f-1) d\underline{u} d\underline{z}$$

where the integration is over the range of values of  $\underline{u}$  and  $\underline{z}$  such that  $\underline{u}'\underline{u} \leq 1$ ,  $\underline{z}'\underline{z} \leq 1$ . Transform from  $\underline{z}$  to  $\underline{\xi} = [\xi_1, \dots, \xi_p]$  by the transformation

$$\underline{\xi} = L\underline{z},$$

where  $L$  is already defined to be a  $p \times p$  orthogonal matrix, whose last row is  $\underline{u}'/\sqrt{\underline{u}'\underline{u}}$ . Then,

$$\underline{z}'\underline{u} = \underline{z}'L'L\underline{u} = \underline{\xi}'L\underline{u} = \xi_p \sqrt{\underline{u}'\underline{u}} = (1-r)^{1/2} \xi_p.$$

Hence (3.9) reduces to

(3.10)  $\int r(1-r)K(\underline{u}|f)d\underline{u} = \int_{-1}^1 K(\underline{t}|f-1)d\underline{t} = K(r-r^2) \cdot \frac{1}{p} K(\underline{t}'\underline{t})$ , due to symmetry of the distribution of  $\underline{t}$ . Now  $\underline{t}$  has the same distribution as  $\underline{u}$  with  $f$  replaced by  $f-1$  and hence finally, (3.10) reduces to

$$\frac{p(f-p+2)}{(f+4)(f+2)} \cdot \frac{1}{f+1}.$$

The covariance between  $r$  and  $s$ , therefore, is (from (3.5))

$$(3.11) \quad \frac{-2p(f-p+2)}{(f+1)(f+2)^2(f+4)}.$$

#### Remarks:

Wilks considers a sample of size  $n$  and has a Wishart matrix based on  $n-1$  degrees of freedom as deviations are from the sample means. He then removes two observations as outliers and thus his  $(n-1)-2$  corresponds to our  $f$ . His  $E(r)$  agrees with our result, with this correspondence but the other moments are in error.

#### Reference

Wilks, S. S. (1963). "Multivariate Statistical Outliers," Sankhyā, Vol. 25, p. 407-426.



UNCLASSIFIED

Security Classification

## DOCUMENT CONTROL DATA - R &amp; D

|   |  |  |                    |
|---|--|--|--------------------|
| 1. REPORT NUMBER<br>SOUTHERN METHODIST UNIVERSITY   |  | 2. REPORT SECURITY CLASSIFICATION<br>UNCLASSIFIED            |                    |
| 3. REPORT DATE<br>August 1, 1970  |  | 4. TOTAL NO OF PAGES<br>6                                    | 5. NO OF REFS<br>1 |
| 6. CONTRACT OR GRANT NO.<br>N00014-68-A-0515<br>PROJECT NO<br>NR042-260   |  | 7. ORIGINATOR'S REPORT NUMBER(S)<br>79                       |                    |
| 8. OTHER REPORT NO. (Any other number that may be assigned this report)   |  |  |                    |
| 9. DISTRIBUTION STATEMENT<br>This document has been approved for public release and sale; its distribution is unlimited. Reproduction in whole or in part is permitted for any purpose of the United States Government.   |  |  |                    |
| 10. SUPPLEMENTARY NOTES   |  | 11. SPONSORING MILITARY ACTIVITY<br>Office of Naval Research |                    |
| 12. ABSTRACT<br>If $B$ is a Wishart matrix and $\underline{x}$ , $\underline{y}$ are two vectors of $p$ components each having a multinormal distribution and if all these quantities are independently distributed, the joint distribution of the two statistics<br>$r = \frac{ B + \underline{y}\underline{y}' }{ B + \underline{xx}' + \underline{y}\underline{y}' } \quad \text{and} \quad s = \frac{ B + \underline{xx}' }{ B + \underline{xx}' + \underline{y}\underline{y}' }$ is derived in this paper. The correlation between $r$ and $s$ is also obtained. $r$ and $s$ are related to Hotelling's $T^2$ and are useful in problems of testing multivariate outliers. |  |  |                    |